

Table 2
Nursing Home Quality Measures (QM)

	Facility A*		Facility B**		Facility C***	
	%		%		%	
	240 (95)		180 (68)		98 (96)	
Ownership	Nonprofit		Government		For Profit	
	2004	2005	2004	2005	2004	2005
High-risk residents who have pressure sores	14	4	13	5	20	9
Low-risk residents who have pressure sores	0	0	4	6	0	0
Residents who spend most of their time in bed or in a chair	3	2	6	8	1	3
Residents with a urinary tract infection	15	21	10	9	5	4
Residents who are more depressed or anxious	8	21	40	22	14	20
Low-risk residents who lose control of their bowels or bladder	75	80	30	40	51	75
Residents whose need for help with daily activities has increased	19	21	27	17	15	19
Residents whose ability to move about in and around their room got worse	9	22	22	25	23	20
Residents who were physically restrained	12	8	12	12	1	0
Residents who have moderate to severe pain	4	4	7	3	14	6
Residents who have/had a catheter inserted and left in their bladder	1	5	10	7	5	6
Short-stay residents who had moderate to severe pain	28	38	20	25	41	24
Short-stay residents with delirium	5	0	4	2	1	3
Short-stay residents with pressure sores	19	13	31	24	25	11

Source: Centers for Medicare and Medicaid Services (February 2004 and February 2005)

Study collection dates: *March–September 2004; **January–July 2004; ***August 2004–February 2005