

**Table 5**

Residents with Cerebral Vascular Accident (18)

<b>Alerts</b>	<b>Triggers</b>	<b>Active Triggers (4380)</b>	
Dehydrated	Left 25% or more of food uneaten.	1261	28.79%
	Dehydrated. Output exceeds input.	257	5.87%
	Diarrhea.	18	0.41%
Skin integrity	Bladder incontinent.	945	21.58%
	Turning or repositioning program has not occurred (within last 6 hours).	165	3.77%
	Bowel incontinent.	148	3.38%
Incontinence	Regular bowel movement has not occurred (within last 72 hours).	158	3.61%
Improvement in condition	Bladder continence decreased from 3 to 2	121	2.76%
	Locomotion off unit decreased from 4 to 3	60	1.37%
	Eating decreased from 1 to 0	57	1.30%
	Locomotion on unit decreased from 4 to 3	54	1.23%
	Walk in corridor decreased from 8 to 3	52	1.19%
	Walk in room decreased from 8 to 3	48	1.10%
	Decision making decreased from 2 to 0	44	1.00%
	Bladder continence decreased from 4 to 2	41	0.94%
	Walk in corridor decreased from 1 to 0	25	0.57%
Decline in condition	Sad or anxious increased from 0 to 2	42	0.96%
	Decisions of daily life severely impaired.	33	0.75%
	Locomotion off unit increased from 1 to 3	26	0.59%
	Bowel continence increased from 2 to 3	23	0.53%