

Table 6

Residents with Dementia (20)

Alerts	Triggers	Active Triggers (9263)	
Skin integrity	Bladder incontinent.	2686	29.00%
	Bowel incontinent.	351	3.79%
	Turning or repositioning program has not occurred (within last 6 hours).	344	3.71%
Dehydration	Left 25% or more of food uneaten.	2233	24.11%
	Dehydrated. Output exceeds input.	299	3.23%
	Diarrhea.	48	0.52%
Improvement in condition	Bladder continence decreased from 3 to 2	254	2.74%
	Locomotion off unit decreased from 4 to 3	176	1.90%
	Bladder continence decreased from 4 to 2	117	1.26%
	Decision making decreased from 2 to 1	112	1.21%
	Decision making decreased from 2 to 0	98	1.06%
	Decision making decreased from 3 to 0	96	1.04%
	Walk in corridor decreased from 8 to 3	92	0.99%
	Locomotion on unit decreased from 1 to 0	60	0.65%
	Eating decreased from 1 to 0	52	0.56%
	Walk in corridor decreased from 1 to 0	46	0.50%
Decline in condition	Toilet use decreased from 3 to 2	46	0.50%
	Decisions of daily life severely impaired.	66	0.71%
Decline in condition	Locomotion off unit increased from 1 to 3	58	0.63%
	Incontinence	Regular bowel movement has not occurred (within last 72 hours).	49