

Table 7
Residents with Depressive Disorder (7)

Alerts	Triggers	Active Triggers (2379)	
Dehydration	Left 25% or more of food uneaten.	733	30.81%
	Dehydrated. Output exceeds input.	99	4.16%
Skin integrity	Bladder incontinent.	620	26.06%
	Turning or repositioning program has not occurred (within last 6 hours).	86	3.61%
	Bowel incontinent.	69	2.90%
Improvement in condition	Decision making decreased from 2 to 0	53	2.23%
	Bladder continence decreased from 3 to 2	45	1.89%
	Walk in corridor decreased from 8 to 3	44	1.85%
	Bladder continence decreased from 4 to 2	29	1.22%
	Personal hygiene decreased from 3 to 2	25	1.05%
	Walk in room decreased from 8 to 3	21	0.88%
	Dressing decreased from 3 to 2	19	0.80%
	Locomotion off unit decreased from 4 to 3	16	0.67%
	Decision making decreased from 2 to 1	15	0.63%
	Locomotion on unit decreased from 1 to 0	14	0.59%
	Bowel continence decreased from 3 to 2	14	0.59%
	Locomotion off unit decreased from 4 to 2	13	0.55%
Decision making decreased from 3 to 0	12	0.50%	
Incontinence	Regular bowel movement has not occurred (within last 72 hours).	50	2.10%
Decline in condition	Dressing increased from 2 to 3	16	0.67%