

Table 8
Residents with Hypertension (11)

Alerts	Triggers	Active Triggers (4644)	
Dehydration	Left 25% or more of food uneaten.	1298	27.95%
	Dehydrated. Output exceeds input.	183	3.94%
Skin integrity	Bladder incontinent.	834	17.96%
	Turning or repositioning program has not occurred (within last 6 hours).	132	2.84%
	Bowel incontinent.	68	1.46%
Improvement in condition	Bladder continence decreased from 3 to 2	132	2.84%
	Decision making decreased from 2 to 0	103	2.22%
	Locomotion off unit decreased from 4 to 3	89	1.92%
	Walk in corridor decreased from 8 to 3	69	1.49%
	Eating decreased from 1 to 0	59	1.27%
	Bladder continence decreased from 4 to 2	56	1.21%
	Walk in room decreased from 8 to 3	56	1.21%
	Locomotion on unit decreased from 4 to 3	53	1.14%
	Transfer decreased from 4 to 3	52	1.12%
	Toilet use decreased from 1 to 0	35	0.75%
	Decision making decreased from 2 to 1	31	0.67%
	Behavioral symptoms decreased from 1 to 0	31	0.67%
	Personal hygiene decreased from 4 to 3	28	0.60%
	Dressing decreased from 4 to 3	28	0.60%
Incontinence	Regular bowel movement has not occurred (within last 72 hours).	66	1.42%