

Table 9

Residents with Osteoarthritis (7)

Alerts	Triggers	Active Triggers (3399)	
Skin integrity	Bladder incontinent.	914	26.89%
	Turning or repositioning program has not occurred (within last 6 hours).	110	3.24%
	Bowel incontinent.	56	1.65%
Dehydration	Left 25% or more of food uneaten.	745	21.92%
	Dehydrated. Output exceeds input.	126	3.71%
Improvement in condition	Bladder continence decreased from 3 to 2	104	3.06%
	Locomotion off unit decreased from 4 to 3	81	2.38%
	Walk in corridor decreased from 8 to 3	67	1.97%
	Walk in room decreased from 8 to 3	63	1.85%
	Bladder continence decreased from 4 to 2	55	1.62%
	Decision making decreased from 3 to 0	42	1.24%
	Locomotion on unit decreased from 4 to 3	29	0.85%
	Personal hygiene decreased from 4 to 3	28	0.82%
	Bed mobility decreased from 4 to 3	28	0.82%
	Toilet use decreased from 4 to 3	28	0.82%
	Eating decreased from 4 to 3	28	0.82%
Decision making decreased from 2 to 1	25	0.74%	
Decline in condition	Sad or anxious increased from 1 to 2	47	1.38%
	Bowel continence increased from 1 to 3	35	1.03%
	Overall condition changed from 2 to 2	30	0.88%